

4-WEBS IN THE PLANE AND THEIR LINEARIZABILITY

Vladislav V. Goldberg

Abstract

We investigate the linearizability problem for different classes of 4-webs in the plane. In particular, we apply a recently found in [AGL] the linearizability conditions for 4-webs in the plane to confirm that a 4-web MW (Mayrhofer's web) with equal curvature forms of its 3-subwebs and a nonconstant basic invariant is always linearizable (this result was first obtained in [M 28]); it also follows from the papers [Na 96] and [Na 98]). Using the same conditions, we also prove that such a 4-web with a constant basic invariant (Nakai's web) is linearizable if and only if it is parallelizable. We also study four classes of the so-called almost parallelizable 4-webs APW_a , $a = 1, 2, 3, 4$ (for them the curvature $K = 0$ and the basic invariant is constant on the leaves of the web foliation X_a), and prove that a 4-web APW_a is linearizable if and only if it coincides with a 4-web MW_a of the corresponding special class of 4-webs MW . The existence theorems are proved for all the classes of 4-webs considered in the paper.

0 Introduction

Let W be a 4-web given by four codimension one foliations X_a , $a = 1, 2, 3, 4$, of curves on a two-dimensional manifold M^2 . The web W is linearizable (rectifiable) if it is equivalent to a linear 4-web, i.e., a 4-web formed by four codimension one foliations of straight lines (see Definition 4 in subsection 2.1).

It is well-known that the geometry of a 4-web W in the plane is completely determined by the curvature K of its 3-subweb $[1, 2, 3]$, defined by its first three foliations X_1, X_2 , and X_3 , and by the basic invariant a and their covariant derivatives (see [G 93], or [G 77], [G 80], [G 88], Section 7.1, where 4-webs of codimension r , $r \geq 1$, on M^{2r} were considered). Note that the basic invariant a is the cross-ratio of the four tangents to the leaves F_2, F_1, F_3, F_4 of the foliations X_2, X_1, X_3, X_4 passing through the point p (see Section 3).

There are many special classes of 4-webs of codimension r , $r > 1$, on M^{2r} and only a few special classes of 4-webs in the plane (of codimension one). The reason for this is that in the case of codimension $r = 1$, most of the classes coincide with the class of parallelizable 4-webs, i.e., 4-webs which are equivalent to a 4-web W formed by four foliations of parallel straight lines of an affine plane $\mathbb{A}^2$ (see Definition 3 in subsection 2.1).

The known (different) special classes of 4-webs in the plane are: parallelizable 4-webs, almost parallelizable 4-webs (4-webs whose basic invariant is constant on the leaves of one of the web foliations), Mayrhofer's 4-webs MW (4-webs with equal curvature forms of their 3-subwebs and nonconstant basic invariant) (see [G 88], Section 7.2, [G 93], and [G 97]) and Nakai's 4-webs NW (4-webs

with a constant basic invariant) (see [Na 96] and [Na 98]). These classes of 4-webs are characterized by some invariant conditions imposed on the curvature K , the basic invariant a , and their covariant derivatives. There is also the class of 4-webs of maximal rank (see [H 94]) but the relations between K , a , and their covariant derivatives for this class are unknown.

A criterion of linearizability is very important in web geometry and in its applications. For 3-webs and 4-webs, the problem of linearizability was posed by Blaschke (see [B 55], §§17 and 42). Blaschke claimed that because of complexity of calculations involving high order differential neighborhoods, it is hopeless to find such a criterion.

Recently the conditions of linearizability for d -webs, $d \geq 4$, on 2-dimensional manifolds were found by Akivis, Goldberg, and Lychagin in [AGL]. For 4-webs, these conditions were presented in two different forms: in the form of two relations expressing the covariant derivatives K_1 and K_2 of the curvature K in terms of the curvature K itself, the basic invariant a and its covariant derivatives a_i, a_{ij}, a_{ijk} of the first three orders, and in the form of two partial differential equations of the fourth order for the web functions $z = f(x, y)$ and $u = g(x, y)$.

In the current paper we consider special classes of 4-webs in the plane and investigate the linearizability problem for them using the first form of linearizability conditions found in [AGL].

There are two different classes of 4-webs with equal curvature forms of its 3-subwebs: 4-webs MW whose basic invariant is nonconstant and 4-webs NW whose basic invariant is constant. We prove the result which follows from the main result of Mayrhofer [M 28] as well as from Nakai's results in [Na 96] and [Na 98]: for 4-webs MW the curvatures and the curvature forms of all their 3-subwebs vanish, i.e., all 3-subwebs are hexagonal (see also [G 93] and [G 97]). 4-webs all 3-subwebs of which are hexagonal were introduced by Mayrhofer (see [M 28]). We arrived to Mayrhofer's 4-webs MW solving the problem of finding 4-webs with a nonconstant basic invariant and equal curvature forms of its 3-subwebs. Mayrhofer was first who proved that such 4-webs are equivalent to 4-webs all foliations of which are pencils of straight lines. There are different proofs of Mayrhofer's theorem: see [M 28], [M 29], [R 28], [BB 38], §10. All these proofs are rather complicated. We think that the reason for this is that the authors of these papers did not use the invariant characterization of 4-webs MW similar to the conditions (42)–(45). Using the linearizability condition of 4-webs indicated above, we present a straightforward short proof of Mayrhofer's theorem (Theorem 10): *a 4-web MW with a nonconstant basic invariant and equal curvature forms of its 3-subwebs is linearizable.*

Note that Mayrhofer's result on 4-webs MW is stronger. He proved that the 4-webs MW are equivalent to 4-webs formed by four pencils of straight lines. However, since in this paper we deal only with 4-web's linearizability, we do not consider these four pencils.

As to the 4-webs with a constant basic invariant and equal curvature forms of its 3-subwebs and (they were studied by Nakai in [Na 96] and [Na 98]), using the same linearizability condition, we prove that *such 4-webs are linearizable if*

and only if they are parallelizable.

We also consider four classes of the so-called almost parallelizable 4-webs APW_a , $a = 1, 2, 3, 4$ (for them the curvature $K = 0$ and the basic invariant is constant on the leaves of one of the web foliations X_a). Using again the same linearizability condition, we prove that *a 4-web APW_a is linearizable if and only if it coincides with a 4-web MW_a of the corresponding special class of 4-webs MW .*

The existence theorems are proved for all classes of 4-webs considered in the paper. Note that some of the classes of 4-webs, for example, almost parallelizable 4-webs APW_a were not known earlier.

1 Basic Notions and Equations

1. 4-webs in the plane. Let M^2 be a two-dimensional analytic manifold.

Definition 1. A 4-web W is given in an open domain D of M^2 by four foliations X_a , $a = 1, 2, 3, 4$, of curves if the tangent lines to these curves through any point $p \in D$ are in general position, i.e., the tangents to the curves of different foliations are distinct and two curves of different foliations have in D at most one common point.

All functions and differential forms which we will introduce on M^2 will be real of class C^∞ , or holomorphic in the complex case.

Definition 2. Two 4-webs W and $\tilde{W}$ with domains $D \subset M^2$ and $\tilde{D} \subset \tilde{M}^2$ are *equivalent* if there exists a diffeomorphism $\phi : D \rightarrow \tilde{D}$ (respectively an analytic isomorphism) of the web domains such that $\phi(X_a) = \tilde{X}_a$.

2. Basic equations of a 3-web. Suppose that (x, y) are local coordinates in $D \subset M^2$ and the foliations X_a , $a = 1, 2, 3, 4$, of a 4-web W are given by the equations

$$u_a(x, y) = \text{const}, \quad a = 1, 2, 3, 4, \quad (1)$$

where $u_a(x, y)$ are differentiable functions.

In order to find the basic equations of a 4-web W , we first consider its 3-subweb $[1, 2, 3]$ defined by the foliations X_α , $\alpha = 1, 2, 3$. If we multiply each of the total differentials du_α of the functions $u_\alpha(x, y)$ from (1) by a factor $g_\alpha(x, y) \neq 0$, we obtain the Pfaffian forms

$$\omega_\alpha = g_\alpha du_\alpha, \quad \alpha = 1, 2, 3. \quad (2)$$

The differential equation $\omega_\alpha = 0$ defines the foliation X_α of curves. The factors g_α can be chosen in such a way that for all the points $(x, y) \in D$ and for all the directions dx/dy , we have the equation

$$\omega_1 + \omega_2 + \omega_3 = 0 \quad (3)$$

(see [B 55], §6).

It follows from (2) that

$$d\omega_\alpha = \omega_\alpha \wedge \theta_\alpha, \quad (4)$$

where

$$\theta_\alpha = -d(\ln g_\alpha).$$

Taking the exterior derivatives of (3) and using (3) and (4), we obtain

$$\omega_1 \wedge (\theta_1 - \theta_3) + \omega_2 \wedge (\theta_2 - \theta_3) = 0.$$

Application of Cartan's lemma to this equation implies

$$\begin{cases} \theta_1 - \theta_3 = a\omega_1 + b\omega_2, \\ \theta_2 - \theta_3 = b\omega_1 + c\omega_2. \end{cases} \quad (5)$$

It follows from (5) that

$$\theta_1 - (a - b)\omega_1 = \theta_3 + b(\omega_1 + \omega_2) = \theta_2 - (c - b)\omega_2 = \theta$$

or

$$\begin{cases} \theta_1 = \theta + (a - b)\omega_1, \\ \theta_2 = \theta + (c - b)\omega_2, \\ \theta_3 = \theta + b\omega_3. \end{cases} \quad (6)$$

Equations (6) allow us to write equations (4) in the form

$$d\omega_\alpha = \omega_\alpha \wedge \theta, \quad \alpha = 1, 2, 3. \quad (7)$$

The form θ is called the *connection form* of the three-web [1, 2, 3].

Exterior differentiation of (7) leads to the cubic exterior equation $d\theta \wedge \omega_\alpha = 0$ from which we easily conclude that

$$d\theta = K\Omega, \quad (8)$$

where K is the *curvature* and

$$\Omega = \omega_1 \wedge \omega_2 = \omega_2 \wedge \omega_3 = \omega_3 \wedge \omega_1 \quad (9)$$

is the *surface element* of the three-web [1, 2, 3]. The form $\Theta = K\Omega$ is called the *curvature form* of the three-web [1, 2, 3]. The equations (8) and (9) are the *structure equations* of the 3-subweb [1, 2, 3] (see [B 55], §8). Equations (7) and (8) prove that the *affine connection* γ_{123} is associated with 3-subweb [1, 2, 3], and θ and K are the *connection form* and the *curvature* of this connection.

3. Basic equations of a 4-web. The fourth foliation of curves of a 4-web W can be given by the equation $\omega_4 = 0$, where

$$-\omega_4 = a\omega_1 + \omega_2 = 0 \quad (10)$$

(see [G 88], Section 7.1, or [G 93]).

The quantity a in equation (10) is called the *basic invariant* of the 4-web W . It satisfies the conditions

$$a \neq 0, 1 \quad (11)$$

(see [G 93], or [G 77], [G 80], [G 88], Section 7.1).

Equations $\omega_a = 0$, $a = 1, 2, 3, 4$, defining the foliations X_a , are preserved under the following concordant transformations of the form ω_a :

$${}'\omega_a = s\omega_a,$$

and only under such transformations. Under such transformations the basic invariant a is not changed, $'a = a$, i.e., a is an absolute invariant.

It is easy to show (see [G 93], or [G 77], [G 80], [G 88], Section 7.1) that *at a point $p \in M^2$, the basic invariant a of the web W equals the cross-ratio of the four tangents to the leaves F_2, F_1, F_3, F_4 of the foliations X_2, X_1, X_3, X_4 passing through the point p .*

The basic invariant a of the 4-web W is an absolute invariant of W . It satisfies the differential equation

$$da = a_1 \omega_1 + a_2 \omega_2, \quad (12)$$

where $a_i \in C^\infty(M^2)$, $i = 1, 2$ are the first covariant derivatives of a .

4. Prolongations of the basic equations. In what follows, we will need prolongations of equations (8). Exterior differentiation of (8) by means of (7) gives the following exterior cubic equation:

$$\nabla K \wedge \omega_1 \wedge \omega_2 = 0, \quad (13)$$

where $\nabla K = dK - 2K\theta$. It follows from (13) that

$$\nabla K = K_1 \omega_1 + K_2 \omega_2, \quad (14)$$

where $K_i \in C^\infty(M^2)$, $i = 1, 2$. Equation (14) shows that the curvature K is a relative invariant of weight two.

We recall that a quantity u is a relative invariant of weight k if

$$du - k\theta = u_1 \omega_1 + u_2 \omega_2$$

(see [AG 96], p. 36) If $k = 0$, then u is an absolute invariant.

We will also need two prolongations of equations (12). Taking exterior derivatives of equations (12), we obtain the following exterior quadratic equation:

$$\nabla a_1 \wedge \omega_1 + \nabla a_2 \wedge \omega_2 = 0, \quad (15)$$

where $\nabla a_i = da_i - a_i \theta$, $i = 1, 2$. Applying Cartan's lemma to equation (15), we find that

$$\nabla a_1 = a_{11} \omega_1 + a_{12} \omega_2, \quad \nabla a_2 = a_{12} \omega_1 + a_{22} \omega_2, \quad (16)$$

where $a_{ij} \in C^\infty(M^2)$, $i, j = 1, 2$.

Taking exterior derivatives of equations (16), we obtain the following exterior quadratic equations:

$$\begin{cases} \nabla a_{11} \wedge \omega_1 + \nabla a_{12} \wedge \omega_2 + a_1 K \omega_1 \wedge \omega_2 = 0, \\ \nabla a_{12} \wedge \omega_1 + \nabla a_{22} \wedge \omega_2 + a_2 K \omega_1 \wedge \omega_2 = 0, \end{cases} \quad (17)$$

where $\nabla a_{ij} = da_{ij} - 2a_{ij}\theta$, $i, j = 1, 2$. Applying Cartan's lemma to equations (17), we find that

$$\begin{cases} \nabla a_{11} = a_{111}\omega_1 + (a_{112} + \frac{2}{3}a_1 K)\omega_2, \\ \nabla a_{12} = (a_{112} - \frac{1}{3}a_1 K)\omega_1 + (a_{122} + \frac{1}{3}a_2 K)\omega_2, \\ \nabla a_{22} = (a_{122} - \frac{2}{3}a_2 K)\omega_1 + a_{222}\omega_2, \end{cases} \quad (18)$$

where $a_{ijk} \in C^\infty(M^2)$, $i, j, k = 1, 2$.

It follows from (16) and (18) that a_i and a_{ij} are relative invariants of weight one and two, respectively.

5. Connection forms, curvatures, and curvature forms of 3-subwebs of a 4-web. Next we will indicate the formulas for the connection forms, the curvatures, and the curvature forms of all 3-subwebs $[1, 2, 3]$ $[1, 2, 4]$, $[1, 3, 4]$ and $[2, 3, 4]$ of a 4-web W .

For the 3-subweb $[1, 2, 3]$, the indicated quantities are:

$$\theta_{123} = \theta, \quad K_{123} = K, \quad \Theta_{123} = K\omega_1 \wedge \omega_2. \quad (19)$$

As was shown in [G 93] (see also [G 88], Section 7.1), for the 3-subweb $[1, 2, 4]$ these quantities are

$$\begin{aligned} \theta_{124} &= \theta - \frac{a_2}{a}\omega_2, \\ K_{124} &= \frac{1}{a} \left(K - \frac{a_{12}}{a} + \frac{a_1 a_2}{a^2} \right), \\ \Theta_{124} &= a K_{124} \omega_1 \wedge \omega_2; \end{aligned} \quad (20)$$

for the 3-subweb $[1, 3, 4]$, they are:

$$\begin{aligned} \theta_{134} &= \theta + \frac{a_2}{1-a}(\omega_1 + \omega_2), \\ K_{134} &= \frac{1}{a-1} \left[K + \frac{a_2(a_1 - a_2)}{(1-a)^2} + \frac{a_{12} - a_{22}}{1-a} \right], \\ \Theta_{134} &= (a-1) K_{134} \omega_1 \wedge \omega_2; \end{aligned} \quad (21)$$

and for the 3-subweb $[2, 3, 4]$, they are

$$\begin{aligned}\theta_{234} &= \theta + \frac{a_1 - a_2}{a} \omega_2 + \frac{a_1}{1-a} (\omega_1 + \omega_2), \\ K_{234} &= \frac{1}{a(a-1)} \left[K + \frac{(2a-1)a_1(a_1-a_2)}{a^2(1-a)^2} + \frac{a_{11} - a_{12}}{a(1-a)} \right], \\ \Theta_{234} &= a(a-1) K_{234} \omega_1 \wedge \omega_2.\end{aligned}\tag{22}$$

2 Special Classes of 4-Webs in the Plane

1. Parallelizable and linearizable 4-webs.

Definition 3. A 4-web W formed by four foliations of parallel straight lines in an affine plane $\mathbb{A}^2$ is said to be a *parallel 4-web*. A 4-web which is equivalent to the parallel 4-web is called *parallelizable*.

For a 3-web in M^2 the definitions of parallel and parallelizable 3-webs are similar.

Remark. A parallel 4-web defined by the foliations X_b , $b = 1, 2, 3, 4$, can be given by the functions

$$u_1 = x, \quad u_2 = y, \quad u_3 = x + y, \quad u_4 = ax + y$$

(cf. [G 93] for parallel 3-webs).

Theorem 1. A 4-web W is parallelizable if and only if the curvature K vanishes, and the basic invariant is constant in M^2 , i.e., if and only if the following conditions are satisfied:

$$K = 0, \quad a_1 = 0, \quad a_2 = 0.\tag{23}$$

Proof. The condition $K = 0$ implies that the leaves of the foliations X_1, X_2 , and X_3 can be defined as level sets of the functions u_1, u_2 , and u_3 in Remark above. The conditions $a_1 = 0, a_2 = 0$ imply that the basic invariant a is constant. Hence leaves of the foliation X_4 are parallel straight lines which are level sets of the function u_4 in Remark. The converse is obvious. $\square$

For another proof see [G 77] or [G 80], or [G 88], Section 7.2.

Definition 4. A 4-web W formed by 4 foliations of straight lines is said to be *linear*. A 4-web which is equivalent to a linear 4-web is called *linearizable* (*rectifiable*).

It follows from Definitions 3 and 4 that a parallelizable 4-web is linearizable but the converse is not true.

In [AGL] the following criterion of linearizability for 4-webs in the plane was found.

Theorem 2. *A 4-web W is linearizable if and only if the curvature K , its covariant derivatives K_1 and K_2 , the basic invariant a , and its covariant derivatives a_i, a_{ij}, a_{ijk} of the first three orders satisfy the following equations:*

$$\begin{aligned}
K_1 = & \frac{1}{a-a^2} \left[\frac{1}{3} (a_1(1-a) + aa_2) K - a_{111} + (2+a)a_{112} - 2aa_{122} \right] \\
& + \frac{1}{(a-a^2)^2} \left\{ \left[(4-6a)a_1 + (-2+3a+a^2)a_2 \right] a_{11} \right. \\
& + \left[(-6+7a+2a^2)a_1 + (2a-3a^2)a_2 \right] a_{12} + \left[(2a(1-a)a_1 - 2a^2a_2) \right] a_{22} \Big\} \\
& + \frac{1}{(a-a^2)^3} \left\{ (-3+8a-6a^2)(a_1)^3 - 2a^3(a_2)^3 \right. \\
& \left. + (6-15a+9a^2+2a^3)(a_1)^2a_2 + (-2a+6a^2-3a^3)a_1(a_2)^2 \right\}, \tag{24}
\end{aligned}$$

$$\begin{aligned}
K_2 = & \frac{1}{a-a^2} \left[\frac{1}{3} (a_1 + a_2(a-1)) K + 2a_{112} - (2a+1)a_{122} + aa_{222} \right] \\
& + \frac{1}{(a-a^2)^2} \left\{ \left[(2a_1 + (2a-2)a_2) \right] a_{11} + \left[(-5+6a)a_1 + (2-3a-2a^2)a_2 \right] a_{12} \right. \\
& + \left[(1-a-2a^2)a_1 + 2a^2a_2 \right] a_{22} \Big\} + \frac{1}{(a-a^2)^3} \left\{ (4a-2)(a_1)^3 + a^3(a_2)^3 \right. \\
& \left. + (5-12a+6a^2)(a_1)^2a_2 + (-2+5a-3a^2-2a^3)a_1(a_2)^2 \right\}. \tag{25}
\end{aligned}$$

Proposition 3. *Linearizable 4-webs exist, and the general solution of the system of equations defining such webs depends on four arbitrary functions of one variable.*

Proof. Taking exterior derivatives of (18) and applying (4), (8), (12), (14), (16), (18), (24), and (25), we get three exterior quadratic equations

$$\begin{aligned}
\nabla a_{111} \wedge \omega_1 + \nabla a_{112} \wedge \omega_2 + (\dots)\omega_1 \wedge \omega_2 &= 0, \\
\nabla a_{112} \wedge \omega_1 + \nabla a_{122} \wedge \omega_2 + (\dots)\omega_1 \wedge \omega_2 &= 0, \\
\nabla a_{122} \wedge \omega_1 + \nabla a_{222} \wedge \omega_2 + (\dots)\omega_1 \wedge \omega_2 &= 0,
\end{aligned} \tag{26}$$

where the coefficients $(\dots)$ depend on K, a, a_i , and a_{ij} . If we substitute the expressions for K_1 and K_2 from (24) and (25) into equation (14) and take exterior derivatives of the resulting equation using the same (4), (8), (12), (14), (16), (18), (24), and (25), we get one more exterior quadratic equation of the

form

$$\begin{aligned} & [-\nabla a_{111} + (2+a)\nabla a_{112} - 2a\nabla a_{122}] \wedge \omega_1 \\ & + [2\nabla a_{112} - (2a+1)\nabla a_{122} + a\nabla a_{222}] \wedge \omega_2 + (\dots)\omega_1 \wedge \omega_2 = 0, \end{aligned} \quad (27)$$

where the coefficients $(\dots)$ again depend on K, a, a_i , and a_{ij} .

The system defining linearizable 4-webs consists of the Pfaffian equation (14) (in which K_1 and K_2 are replaced by their values (24) and (25)), (3), (16), (18), and exterior quadratic equations (26), (27). For this system we have 4 unknown forms ∇a_{ijk} . Hence $q = 4$. Since the exterior quadratic equations (26) and (27) are obviously independent, and their number is four, we have $s_1 = 4$. Thus $s_2 = q - s_1 = 0$, and the Cartan number $Q = s_1 + 2s_2 = 4$. On the other hand, if we apply Cartan's lemma to (26), we find the forms ∇a_{ijk} in terms of ω_i , and there will be 5 coefficients in the expansions of these forms. Substituting these expansions into (27), we find one of these coefficients. So, the dimension N of the space of integral elements over a point equals four, $N = 4$. Thus $Q = N$, and by Cartan's test (see [BCGGG 91], p. 120), the system is in involution, and its general solution depends on four functions of one variable. $\square$

2. 4-webs whose basic invariant is constant on the leaves of one of web foliations.

Theorem 4. *The basic invariant a of a 4-web is constant on the leaves of the web foliation X_1, X_2, X_3 , and X_4 if and only if there is respectively the following relation between the covariant derivatives a_1 and a_2 of this invariant:*

$$a_2 = 0, \quad (28)$$

$$a_1 = 0, \quad (29)$$

$$a_1 = a_2, \quad (30)$$

$$a_1 = aa_2. \quad (31)$$

Proof. In fact, it follows from (1.3) that conditions (29)–(31) are necessary and sufficient for the following congruences:

$$da \equiv 0 \pmod{\omega_1},$$

$$da \equiv 0 \pmod{\omega_2},$$

$$da \equiv 0 \pmod{\omega_1 + \omega_2},$$

$$da \equiv 0 \pmod{a\omega_1 + \omega_2},$$

respectively. $\square$

Proposition 5. *4-webs whose basic invariant is constant on the leaves of one of web foliations exist, and the general solution of the system of equations defining such webs depends on one arbitrary function of two variables.*

Proof. Suppose, for example, that the basic invariant is constant on the leaves of the foliation X_2 defined by the equation $\omega_2 = 0$. Then condition (29) holds. For such a 4-web, we have the Pfaffian equations (12) and (16). By (29), the former becomes

$$da = a_2\omega_2. \quad (32)$$

Taking the exterior derivatives of these two Pfaffian equations, we arrive at the following two exterior quadratic equations:

$$\begin{aligned} \nabla K_1 \wedge \omega_1 + \nabla K_2 \wedge \omega_2 &= 0, \\ \nabla a_2 \wedge \omega_2 &= 0, \end{aligned} \quad (33)$$

where $\nabla K_i = dK_i - 3K_i\theta$, $\nabla a_2 = da_2 - a_2\theta$. Thus, we have three unknown functions, and $q = 3$. Since there are two exterior quadratic equations, we have $s_1 = 2$. As a result, $s_2 = q - s_1 = 1$, and the Cartan number $Q = s_1 + 2s_2 = 4$. Applying Cartan's lemma to the exterior quadratic equations (33), we find that

$$\begin{aligned} \nabla K_1 &= K_{11}\omega_1 + K_{12}\omega_2, \\ \nabla K_2 &= K_{12}\omega_1 + K_{22}\omega_2, \\ \nabla a_2 &= a_{22}\omega_2. \end{aligned}$$

Hence the dimension N of the space of integral elements over a point equals 4, $N = 4$. Therefore, $Q = N$, and by Cartan's test, our system of equations is in involution, and its general solution depends on one arbitrary function of two variables.

The proof in other cases is similar. The difference is that in the cases when a 4-web satisfies equations (28), (30), and (31), equation (32) has respectively the form

$$\begin{aligned} da &= a_1\omega_1, \\ da &= a_1(\omega_1 + \omega_2), \end{aligned}$$

and

$$da = a_2(a\omega_1 + \omega_2),$$

and the last equation (33) becomes

$$\begin{aligned} \nabla a_1 \wedge \omega_1 &= 0, \\ \nabla a_1 \wedge (\omega_1 + \omega_2) &= 0, \end{aligned}$$

and

$$\nabla a_2 \wedge (a\omega_1 + \omega_2) = 0,$$

respectively. □

3. Almost parallelizable 4-webs.

Definition 5. We say that a 4-web W is *almost parallelizable* if its 3-subweb $[1, 2, 3]$ is parallelizable (i.e., $K = 0$), and its basic invariant is constant on the leaves of one of the web foliations.

So, according to Theorem 4, there are four classes of almost parallelizable 4-webs characterized by the following conditions:

$$K = 0, \quad a_2 = 0, \quad (34)$$

$$K = 0, \quad a_1 = 0, \quad (35)$$

$$K = 0, \quad a_1 = a_2, \quad (36)$$

$$K = 0, \quad a_1 = aa_2. \quad (37)$$

We denote the 4-webs of these four classes by APW_a , $a = 1, 2, 3, 4$.

Proposition 6. *The almost parallelizable 4-webs APW_a exist, and the general solution of the system of equations defining such webs depends on one arbitrary function of one variable.*

Proof. In fact, in each of the cases (34)–(37), the condition $K = 0$ implies $K_1 = 0$, $K_2 = 0$. Thus, we have $q = 1$. For a 4-web APW_1 , in (33) (and in similar systems corresponding to 4-webs APW_2 , APW_3 , and APW_4) only the second equation remains. Hence Cartan's characters are $s_1 = 1$ and $s_2 = 0$. It is easy to see that in this case $Q = N = 1$, and by Cartan's test, our system of equations is involutive, and its general solution depends on one arbitrary function of one variable. $\square$

The following theorem establishes the values of the curvatures K_{abc} for the 4-webs APW_a .

Theorem 7. *The curvatures K_{abc} for the 4-webs APW_a are*

(i) *For the 4-webs APW_1 :*

$$\frac{K}{123} = \frac{K}{124} = \frac{K}{134} = 0, \quad \frac{K}{234} = \frac{(1-2a)a_1^2 - (a-a^2)a_{11}}{(a-a^2)^3}; \quad (38)$$

(ii) *For the 4-webs APW_2 :*

$$\frac{K}{123} = \frac{K}{124} = \frac{K}{234} = 0, \quad \frac{K}{134} = \frac{a_2^2 + (1-a)a_{22}}{(1-a)^3}; \quad (39)$$

(iii) For the 4-webs APW_3 :

$$K_{123} = K_{134} = K_{234} = 0, \quad K_{124} = \frac{a_1^2 - aa_{11}}{a^3}; \quad (40)$$

(iv) For the 4-webs APW_4 :

$$K_{123} = 0, \quad K_{124} = -\frac{aa_{22}}{a^2}, \quad K_{134} = \frac{a_{22}}{1-a}, \quad K_{234} = \frac{a_{22}}{a-a^2}. \quad (41)$$

Proof. The proof is straightforward: it follows from formulas (20)–(22) and conditions (34)–(37). $\square$

4. 4-webs with a nonconstant basic invariant and equal curvature forms of their 3-subwebs. It is important to emphasize that we assume here that *the basic invariant of a 4-web is not constant on M_2* , i.e., its covariant derivatives a_1 and a_2 do not vanish simultaneously. This assumption shows that the class of 4-webs we are going to consider is completely different from the class considered by Nakai in [Na 96] and [Na 98]. Nakai also considered 4-webs with equal curvature forms of their 3-subwebs. Assuming that these equal curvature forms of the 3-subwebs are nonvanishing, he proved that the basic invariant a of a 4-web is constant on M_2 , i.e., $a_1 = a_2 = 0$. Thus in [Na 96] and [Na 98], Nakai considered 4-webs with a constant basic invariant. We will consider this kind of 4-webs in subsection 6.

Theorem 8. *A 4-web W has a nonconstant basic invariant a and equal curvature forms of its 3-subwebs $[a, b, c]$ if and only if all 3-subwebs $[a, b, c]$ are parallelizable, i.e., if*

$$K_{abc} = 0, \quad \Theta_{abc} = 0, \quad a, b, c = 1, 2, 3, 4, \quad (42)$$

and the second covariant derivatives a_{ij} of the basic invariant a are expressed in terms of the invariant a itself and its first covariant derivatives a_i as follows:

$$a_{11} = \frac{a_1[(1-2a)a_1 + aa_2]}{a-a^2}, \quad (43)$$

$$a_{12} = \frac{a_1 a_2}{a}, \quad (44)$$

$$a_{22} = \frac{a_2(a_1 - aa_2)}{a-a^2}. \quad (45)$$

Proof. In fact, by equation (20), it follows that if $a \neq 0$, then the condition $\Theta_{124} = \Theta_{123}$ holds if and only if equation (44) is valid.

In a similar manner, by (21) and (22), it follows that if $a \neq 0$, then the conditions $\Theta_{134} = \Theta_{123}$ and $\Theta_{234} = \Theta_{123}$ hold if and only if equations (45) and (43) are valid, respectively.

To prove (42), we substitute the values of a_{ij} from (43)–(45) into equations (16). This gives

$$\begin{aligned} da_1 &= a_1\theta + \frac{[(1-2a)a_1 + aa_2]a_1}{a-a^2}\omega_1 + \frac{a_1a_2}{a}\omega_2, \\ da_2 &= a_2\theta + \frac{1}{a-a^2}[a_1a_2(1-a)\omega_1 + a_2(a_1 - aa_2)\omega_2]. \end{aligned} \quad (46)$$

Exterior differentiation of either of equations (46) implies $K = 0$. The latter condition and equations (43)–(45) lead to (42).

Thus, each of 3-subwebs $[a, b, c]$ is parallelizable. However, by Theorem 1, this does not imply the parallelizability of the 4-web W —the latter web is parallelizable if and only if all 3-subwebs $[a, b, c]$ are parallelizable and the basic invariant a is constant in the connection γ_{123} , i.e., if and only if we have conditions (42) and $a_1 = a_2 = 0$. $\square$

We denote the 4-webs with equal curvature forms of their 3-subwebs and a nonconstant basic invariant by MW .

Note that conditions (42) were first obtained by Nakai (see [Na 96] and [Na 98]). As to conditions (43)–(45) characterizing the webs MW , they appeared here at the first time.

4-webs all 3-subwebs of which are parallelizable (hexagonal) were introduced by Mayrhofer (see [M 28]). This is the reason that we denoted them by MW (Mayrhofer's 4-webs). We arrived to Mayrhofer's 4-webs MW solving the problem of finding 4-webs with a nonconstant basic invariant and equal curvature forms of its 3-subwebs. Mayrhofer was first who proved that such 4-webs are equivalent to 4-webs all foliations of which are pencils of straight lines. There are different proofs of Mayrhofer's theorem: see [M 28], [M 29], [R 28], [BB 38], §10. However, all these proofs are rather complicated. We think that the reason for this is that the authors of these papers did not use the invariant characterization of 4-webs MW similar to the conditions (42)–(45).

Now we prove the existence theorem for webs NW .

Proposition 9. *The 4-webs MW exist. The system of Pfaffian equations defining such webs is completely integrable, and its general solution depends on three arbitrary constants.*

Proof. For a web MW , we have Pfaffian equations (12) and (16), where the a_{ij} are expressed by formulas (43)–(45). If we take exterior derivatives of equations (16) in which a_{ij} are replaced by their values (43)–(45), we arrive at the identities. So, for the system in question, we have $s_0 = 3$, $s_1 = 0$. The system is completely integrable, and its general solution depends on three arbitrary constants. $\square$

In the next theorem we present a straightforward short proof of the fact that 4-webs MW are always linearizable by applying a linearizability condition of 4-webs found in [AGL]. This theorem gives the first nontrivial application of linearizability conditions (24) and (25) for 4-webs.

Theorem 10. (Mayrhofer [M 28]) A 4-web MW is linearizable.

Proof. To prove this result, we must prove that conditions (24) and (25) of linearizability are satisfied identically for the 4-webs MW . To check these conditions, we substitute the values of the invariants a_{ij} and a_{ijk} into them. As to a_{ij} , their values are given by (43)–(45).

In order to find a_{ijk} , we differentiate (43)–(45) by using (12) and (16) and compare the result with equations (18). Equating the coefficients of ω_1 and ω_2 in the resulting equations, we find that

$$\begin{aligned} a_{111} &= \frac{1}{(a-a^2)^2} \left[(6a^2-6a+1)a_1^3 + (4a-6a^2)a_1^2a_2 + a^2a_1a_2^2 \right], \\ a_{112} &= \frac{1}{(a-a^2)^2} \left[(1-2a)(1-a)a_1^2a_2 + (a-a^2)a_1a_2^2 \right] = \frac{a_2a_{11}}{a}, \\ a_{122} &= \frac{a_1a_2(a_1-aa_2)}{a^2(1-a)} = \frac{a_1a_{22}}{a}, \\ a_{222} &= \frac{1}{(a-a^2)^2} (a_1^2a - 2aa_1a_2^2 + a^2a_2^3). \end{aligned} \tag{47}$$

Next, by (42), for 4-webs MW , we have $K = 0$. This and (14) imply $K_1 = K_2 = 0$. Substituting the values of a_{ij} from (43)–(45) and the values of a_{ijk} from (47) into (24) and (25), after lengthy straightforward calculations, we come to the identities: the coefficients of a_1^3 , $a_1^2a_2$, $a_1a_2^2$ and a_2^3 vanish. Hence any 4-web MW is linearizable. $\square$

5. Special classes of 4-webs MW . Although for 4-webs MW the basic invariant a is not constant on M^2 , it can be constant on the leaves of one of the foliations X_a of a web MW . If this is the case, we have four special classes of 4-webs MW . These four classes are intersections of four classes of almost parallelizable 4-webs APW_a and the class of 4-webs MW . We denote webs of these classes by MW_a .

Theorem 11. *The 4-webs MW_a are characterized by the following conditions:*

$$\begin{aligned} \text{Webs } MW_1 : \quad & K = 0, \quad a_2 = 0, \quad a_{11} = \frac{(1-2a)a_1^2}{a-a^2}, \quad a_{12} = 0, \quad a_{22} = 0; \\ \text{Webs } MW_2 : \quad & K = 0, \quad a_1 = 0, \quad a_{11} = 0, \quad a_{12} = 0, \quad a_{22} = -\frac{a_2^2}{1-a}; \\ \text{Webs } MW_3 : \quad & K = 0, \quad a_1 = a_2, \quad a_{11} = a_{12} = a_{22} = \frac{a_1^2}{a}; \\ \text{Webs } MW_4 : \quad & K = 0, \quad a_1 = aa_2, \quad a_{11} = 2aa_2^2, \quad a_{12} = a_2^2, \quad a_{22} = 0. \end{aligned} \tag{48}$$

Proof. The proof follows from (34)–(37) and (43)–(45). $\square$

Note that by (48) and (38)–(41), for any 4-web MW_a , all the curvatures K_{abc} vanish (as it should be according to Theorem 8 (see (42))).

Proposition 12. *The 4-webs MW_a , $a = 1, 2, 3, 4$, exist. The system of Pfaffian equations defining such webs is completely integrable, and its general solution depends on two arbitrary constants.*

Proof. In fact, by (48), in the system defining each of the webs MW_a (see proof of Theorem 9), only two Pfaffian equations are independent. $\square$

The next theorem gives a relation between the 4-webs APW_a and MW_a .

Theorem 13. *A 4-web APW_a , $a = 1, 2, 3, 4$, is a 4-web MW_a if and only if it is linearizable.*

Proof. The necessity is obvious: any MW_a is APW_a , and by Theorem 10, any MW_a is linearizable.

We prove the sufficiency separately for different $a = 1, 2, 3, 4$.

- (i) *Linearizable 4-webs APW_1 .* Note that it follows from (34), (16), and (18) that for a 4-web APW_1 not only $a_2 = 0$ but also $a_{12} = a_{22} = a_{112} = a_{122} = a_{222} = 0$. Since by hypothesis our 4-web APW_1 is linearizable, the linearizability conditions (24) and (25) must be satisfied identically. Substituting

$$K = K_1 = K_2 = 0, \quad a_1 = a_{11} = a_{12} = a_{111} = a_{112} = a_{122} = 0$$

into (25) and (24), we find that

$$a_{11} = -\frac{(1-2a)a_1^2}{a-a^2}$$

and

$$a_{111} = -\frac{(6a^2-6a+1)a_1^3}{(a-a^2)^2}.$$

Comparing with (48), we see that a *linearizable 4-web APW_1* is a 4-web MW_1 . As to the second condition obtained from (24), it is easy to see that it is a differential consequence of the first condition.

- (ii) *Linearizable 4-webs APW_2 .* Note that it follows from (34), (16), and (18) that for a 4-web APW_2 not only $a_1 = 0$ but also $a_{11} = a_{12} = a_{111} = a_{112} = a_{122} = 0$. Since by hypothesis, our 4-web APW_2 is linearizable, the linearizability conditions (24) and (25) must be satisfied identically. Substituting

$$K = K_1 = K_2 = 0, \quad a_1 = a_{11} = a_{12} = a_{111} = a_{112} = a_{122} = 0$$

into (24) and (25), we find that

$$a_{22} = -\frac{a_2^2}{1-a}$$

and

$$a_{222} = -\frac{a_2^3}{(1-a)^2}.$$

Comparing with (48), we see that a *linearizable 4-web* APW_2 is a 4-web MW_2 . As to the second condition obtained from (25), it is easy to see that it is a differential consequence of the first condition.

- (iii) *Linearizable 4-webs* APW_3 . Note that it follows from (34), (16), and (18) that for a 4-web APW_3 , the condition $a_1 = a_2$ implies $a_{11} = a_{12} = a_{22}$ and $a_{111} = a_{112} = a_{122} = a_{222}$. Since by hypothesis, our 4-web APW_3 is linearizable, the linearizability conditions (24) and (25) must be satisfied identically. Substituting

$$K = K_1 = K_2 = 0, \quad a_1 = a_2, \quad a_{11} = a_{12} = a_{22}, \quad a_{111} = a_{112} = a_{122} = a_{222}$$

into (24) and (25), we find that

$$a_{111} = \frac{4a_1a_{11}}{a} - \frac{3a_1^3}{a^2}$$

and

$$a_{111} = -\frac{2a_1a_{11}}{a^2} - \frac{a_1^3}{a^3}.$$

It follows from these two equations that

$$a_{11} = -\frac{a_1^2}{a}.$$

Comparing this with (48), we see that a *linearizable 4-web* APW_3 is a 4-web MW_3 . As to the expression for a_{111} , it is easy to see that this expression can be obtained by differentiation from the expression of a_{11} .

- (iv) *Linearizable 4-webs* APW_4 . Note that it follows from (34), (16), and (18) that for a 4-web APW_4 , the condition $a_1 = aa_2$ implies

$$a_{11} = a^2a_{22} + 2a_2^2, \quad a_{12} = aa_{22} + a_2^2$$

and

$$\begin{aligned} a_{111} &= a^3a_{222} + (7a^2a_2 + 2aa_2)a_{22} + 6aa_2^3, \\ a_{112} &= a^2a_{222} + 6aa_2a_{22} + 2a_2^3, \\ a_{122} &= aa_{222} + 3a_2a_{22}. \end{aligned}$$

Since by hypothesis, our 4-web APW_4 is linearizable, the linearizability conditions (24) and (25) must be satisfied identically. Substituting

$$K = K_1 = K_2 = 0, \quad a_1 = aa_2$$

and the expressions for $a_{11}, a_{12}, a_{111}, a_{112}, a_{122}$ written above into (24) and (25), we find that (25) is identically satisfied, and (24) reduces to

$$a_{22} = 0.$$

As a result, we see from the expressions for a_{11} and a_{12} that they become

$$a_{11} = 2aa_2^2, \quad a_{12} = a_2^2.$$

Comparing the results with (48), we see that a *linearizable 4-web* APW_4 is a 4-web MW_4 .

□

The next theorem gives another criterion for a 4-web APW_a to be a 4-web MW_a .

Theorem 14. *A 4-web APW_a , $a = 1, 2, 3, 4$, is a 4-web MW_a if and only if all its 3-subwebs are parallelizable.*

Proof. By Theorem 7, for 4-webs APW_1, APW_2 , and APW_3 , three of the 3-subwebs are parallelizable. It follows from (38)–(40) that the fourth 3-subweb of these 4-webs is parallelizable if and only if we have

$$a_{22} = -\frac{a_2^2}{1-a}, \quad a_{11} = \frac{(1-2a)a_1^2}{a-a^2}, \quad a_{11} = a_{12} = a_{22} = \frac{a_1^2}{a},$$

respectively. Comparing these relations with (48), we see that 4-webs APW_1, APW_2 , and APW_3 are 4-webs MW_1, MW_2 , and MW_3 , respectively.

For a 4-web APW_4 , by (41), only the 3-subweb $[1, 2, 3]$ is parallelizable. But relations (41) show also that other 3-subwebs are parallelizable if and only if

$$a_{22} = 0.$$

But as we already showed in the proof of Theorem 13, the condition $a_{22} = 0$ implies

$$a_{11} = 2aa_2^2, \quad a_{12} = a_2^2,$$

and by (48), our 4-web APW_4 is a 4-web MW_4 .

□

6. 4-webs with a constant basic invariant and equal curvature forms of their 3-subwebs. For such 4-webs, we have

$$a_1 = a_2 = 0. \tag{49}$$

By (14) and (16), conditions (49) imply

$$a_{ij} = 0, \quad a_{ijk} = 0, \quad i, j, k = 1, 2. \tag{50}$$

Now formulas (20)–(22) imply

$$\frac{K}{123} = \frac{K}{124} = \frac{K}{134} = \frac{K}{234} = K$$

and

$$\Theta_{123} = \Theta_{124} = \Theta_{134} = \Theta_{234} = K\omega_1 \wedge \omega_2,$$

i.e., the curvatures and the curvature forms of all 3-subwebs are equal without vanishing.

We denote such 4-webs by *NW* (Nakai's webs—see [Na 96] and [Na 98]).

Proposition 15. *The 4-webs NW exist, and the general solution of the system of equations defining such webs depends on one arbitrary function of two variables.*

Proof. In fact, for 4-webs *NW*, equation (12) becomes $da = 0$. Taking exterior derivatives of (14), we arrive at the exterior quadratic equation

$$\nabla K_1 \wedge \omega_1 + \nabla K_2 \wedge \omega_2 = 0,$$

where $\nabla K_i = dK_i - 3K_i\theta$. It is easy to see that in this case, we have $q = 2$, the Cartan characters $s_1 = 1$, $s_2 = 1$, and $Q = s_1 + 2s_2 = 3$.

Next, it follows from the quadratic equation above that

$$\nabla K_1 = K_{11}\omega_1 + K_{12}\omega_2, \quad \nabla K_2 = K_{12}\omega_1 + K_{22}\omega_2.$$

Hence the dimension N of the space of integral elements over a point equals three, $N = 3$. Therefore, $Q = N$, and by Cartan's test, our system of equations is in involution, and its general solution depends on one arbitrary function of two variables. $\square$

It is easy to see that, in general, a 4-web *NW* is not linearizable since conditions (24) and (25) do not hold for such a web.

However, equations (24) and (25) produce the simple conditions for linearizability of 4-webs *NW*. Namely, by (49) and (50), it follows from (24) and (25) that

$$K_1 = 0, \quad K_2 = 0. \tag{51}$$

Now we are able to prove the following theorem describing linearizable 4-webs *NW*.

Theorem 16. *If a 4-web NW is linearizable, then it is parallelizable.*

Proof. In fact, for a linearizable 4-web *NW*, we have equations (51). They and equation (14) imply

$$dK - 3K\theta = 0.$$

Taking exterior derivative of this equation and applying (8), we find that

$$-2K^2\omega_1 \wedge \omega_2 = 0.$$

It follows that $K = 0$, and by Theorem 1, this and (49) prove that a linearizable 4-web *NW* is parallelizable. $\square$

Theorem 16 implies the following corollaries.

Corollary 17. *There exists no linearizable nonparallelizable 4-webs NW.*

Corollary 18. *Parallelizable 4-webs NW exist. They are defined by one completely integrable equations $da = 0$. The set of such parallelizable 4-webs NW depends on one constant (the constant basic invariant a).*

Proof. In fact, since the 4-web NW is parallelizable, we have $K = K_1 = K_2 = 0$. So, the only remaining equation defining parallelizable 4-webs NW is the equation $da = 0, a \neq 0, 1$. This proves the corollary. $\square$

References

- [AG 96] Akivis, M. A. and V. V. Goldberg, *Conformal Differential Geometry and Its Generalizations*, John Wiley & Sons, 1996, xiv+383 pp. (MR 98a:53023; Zbl. 863.53002.)
- [AGL] Akivis, M. A., V. V. Goldberg, and V. V. Lychagin, *Linearizability of d -webs, $d \geq 4$, on 2-dimensional manifolds*, Preprint, 2002, 17 pp., arXiv: math.DG/0209290.
- [B 55] Blaschke, W., *Einführung in die Geometrie der Waben*, Birkhäuser-Verlag, Basel-Stuttgart, 1955, 108 pp. (MR 17, 780; Zbl. 68, 365); Russian transl., GITTL, Moskva, 1959, 144 pp. (MR 22 #2942.)
- [BB 38] Blaschke, W. and G. Bol, *Geometrie der Gewebe*, Springer-Verlag, Berlin, 1938, viii+339 pp. (MR 6, p. 19; Zbl. 20, p. 67.)
- [BCGGG 91] Bryant, R. L., S. S. Chern, R. B. Gardner, H. L. Goldsmith, and P. A. Griffiths, *Exterior differential systems*, Springer-Verlag, New York, 1991, vii+475 pp. (MR 92h:58007; Zbl. 726.58002.)
- [Ca 23] Cartan, É., *Sur les variétés à connexion affine et la théorie de la relativité généralisée*, Ann. École Norm. **40** (1923), 325–412. (JFM 50, p. 542); *Œuvres complètes: Partie III, Divers, géométrie, différentielle*, vol. 3, Gauthier-Villars, Paris, 1955, pp. 659–746. (MR 17, 697; Zbl. 15, 83.)
- [G 77] Goldberg, V. V., *On the theory of four-webs of multidimensional surfaces on a differentiable manifold X_{2r}* , Izv. Vyssh. Uchebn. Zaved. Mat. **1977**, no. 11(186), 15–22 (Russian), English transl., Soviet Math.(Iz. VUZ) **21** (1977), no. 11, 97–100. (MR 58 #30859; Zbl. 398.53009 & 453.53010.)
- [G 80] Goldberg, V. V., *On the theory of four-webs of multidimensional surfaces on a differentiable manifold X_{2r}* , Serdica **6** (1980), no. 2, 105–119 (Russian). (MR 82f:53023; Zbl. 453.53010.)

- [G 88] Goldberg, V. V., *Theory of multicodimensional $(n + 1)$ -webs*, Kluwer Academic Publishers, Dordrecht, 1988, xxii+466 pp. (MR 89h:53021; Zbl. 668.53001.)
- [G 93] Goldberg, V. V., *Curvilinear 4-webs with equal curvature forms of its 3-subwebs*, Webs and Quasigroups, Tver State Univ., Tver, 1993, 9–19. (MR 94g:53012; Zbl. 792.53010.)
- [G 97] Goldberg, V. V., *Special classes of curvilinear 4-webs with equal curvature forms of its 3-subwebs*, Webs and Quasigroups, Tver State Univ., Tver, 1996–1997, 24–39 (1997). (MR 2001f:53037; Zbl. 903.53013.)
- [H 94] Hénaut, A., *Caractérisation des tissus de C^2 dont le rang est maximal et qui sont linéarisables*, Composito Math. **94** (1994), no. 3, 247–268. (MR 96a:32057; Zbl. 877.53013.)
- [KN 63] Kobayashi, S. and K. Nomizu, *Foundations of differential geometry*, vol. 1, Wiley–Interscience, New York–London–Sydney, 1963, xi+329 pp. (MR **27** #2945; Zbl. **119**, p. 375.)
- [M 28] Mayrhofer, K.: *Kurvensysteme auf Flächen*, Math. Z. **28** (1928), 728–752. (JFM **54**, p. 745.)
- [M 29] Mayrhofer, K.: *Über Sechsensysteme*, Abh. Math. Sem. Univ. Hamburg **7** (1929), 1–10. (JFM **55**, p. 1013.)
- [Na 96] Nakai, I., *Curvature of curvilinear 4-webs and pencils of one forms*, Topology of holomorphic dynamical systems and related topics (Kyoto, 1995), Sūrikaiseikikenkyūsho Kōkyūroku No. 955 (1996), no. 8, 109–132. (MR 98a:53026.)
- [Na 98] Nakai, I., *Curvature of curvilinear 4-webs and pencils of one forms: Variation on the theme a theorem of Poincaré, Mayrhofer and Reidemeister on curvilinear 4-webs*, Comment. Math. Helv. **73** (1998), 177–205. (MR 99c:53013; Zbl. 926.53011.)
- [R 28] Reidemeister, K.: *Gewebe und Gruppen*, Math. Z. **29** (1928), 427–435. (JFM **54**, p. 746.)

Author's address: Department of Mathematical Sciences, New Jersey Institute of Technology, University Heights, Newark, N.J. 07102, U.S.A.

E-mail address: vlgold@m.njit.edu